

# Practice Pearl Article: Avoiding Multidrug Resistance in a Hospital Setting

Grace Johanna Salazar Tamayo<sup>1,2,3</sup>

1. Infection control and prevention, Axxis Hospital, Quito, Ecuador; 2. Infectious disease department, Vozandes Hospital, Quito, Ecuador; 3. Infectious disease department, Metropolitano Hospital, Quito, Ecuador

<https://doi.org/https://doi.org/10.17925/ID.2024.3.1.3>

**A**voiding multidrug resistance in hospitals is a challenging but achievable task that requires understanding local epidemiology and evaluating and intervening in cleaning and disinfection processes, antibiotic stewardship and prevention measures. These activities must be routinely evaluated and continuously applied, with executive directive management support and teamwork from infection committee members alongside active leaders of each process. An infographic on avoiding multidrug resistance in a hospital setting is available alongside this article here.

## Keywords

Antimicrobial stewardship, bacterial infections - prevention & control, cross infection - prevention & control, disinfection, drug resistance, hand hygiene, infection control - methods, infection control - organization & administration

**Disclosures:** Grace Johanna Salazar Tamayo has no financial or non-financial relationships or activities to declare in relation to this article.

**Acknowledgements:** This article is published with an infographic to enhance understanding, which was peer reviewed alongside the manuscript. To view the infographic, please see via [www.touchinfectiousdiseases.com](http://www.touchinfectiousdiseases.com) here.

**Review Process:** Double-blind peer review.

**Compliance with ethics:** This article is an opinion piece and does not report on new clinical data, or any studies with human or animal subjects performed by any of the authors.

**Data availability:** Data sharing is not applicable to this article as no datasets were generated or analysed during the writing of this article.

**Authorship:** The named author meets the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, takes responsibility for the integrity of the work as a whole, and has given final approval for the version to be published.

**Access:** This article is freely accessible at [touchINFECTIOUSDISEASES.com](http://touchINFECTIOUSDISEASES.com). ©Touch Medical Media 2024.

**Received:** 13 December 2023

**Accepted:** 24 June 2024

**Published online:** 25 September 2024

**Citation:** *touchREVIEWS in Infectious Diseases*. 2024;3(1):7–10

**Corresponding author:** Dr Grace Johanna Salazar Tamayo, MD, Infection control and prevention, Axxis Hospital, Quito, Ecuador CP 170401. E: [grace.salazar@gmail.com](mailto:grace.salazar@gmail.com)

**Support:** No funding was received in the publication of this article.

## Highlights

- Determining infection and multidrug resistance (MDR) rates continuously helps to detect the problem to be solved, including outbreaks.
- Cleaning and disinfection play an important role in preventing MDR and evaluating and correcting the process for better results.
- MDR control can only be achieved with various measures applied continuously and consistently by a group of committed individuals.

## Development

Antimicrobial resistance is a global crisis. Approximately 1.3 million deaths were directly attributable to multidrug-resistant (MDR) microorganisms in 2019.<sup>1</sup> These infections also have significant implications for hospital control and expenditure, as well as morbidity and mortality.<sup>2,3</sup> MDR microorganisms can colonize a patient for months or years, and this colonized population can re-enter the community and potentially transmit this resistance.<sup>4,5</sup>

Controlling MDR in hospitals is crucial not only for reducing morbidity and mortality in hospitalized patients, but also for ensuring the quality of healthcare and reducing hospital costs.<sup>6</sup>

This task is challenging and involves multiple activities implemented simultaneously after a local epidemiological analysis, along with evaluating and providing feedback on the measures taken to identify the strengths and weaknesses of each hospital environment. These activities include the following:

1. evaluating the epidemiological profile of community- and healthcare-associated infections (HAIs);
2. assessing the hospital resistance profile;
3. analysing antibiotic consumption;
4. evaluating cleaning and disinfection processes;
5. promoting hand hygiene; and
6. implementing other measures.

## Point 1: Evaluating the epidemiological profile of community- and healthcare-associated infections

This includes identifying the infections, causative microorganisms and susceptibility profiles reported as a value, for example, #/1,000 catheter days and #/1,000 ventilator days, which are evaluated over a specific period and are globally comparable.<sup>7,8</sup> The reported frequency of MDR in HAIs varies between 25.4% and 42.5%, and these percentages vary in each hospital.<sup>9–11</sup> Analysing the rates of infections and the frequency of MDR microorganisms and whether empirical antimicrobial use is appropriate in HAIs benefit the problem detection, including outbreaks. These data must be collected, evaluated and communicated continuously because it is the key to management decision-making.

Table 1: Outbreaks with different control and prevention interventions<sup>29–33</sup>

Study	Problem	Intervention in cleaning and disinfection	Other interventions	Main outcome
Ikeda et al. <sup>29</sup>	Outbreak of ESBL-producing bacteria in catheter-associated urinary tract infection in patients with spinal cord injury	Cleaning procedures were reviewed and the number of routine cleanings was increased using a complex-chlorine-based disinfectant cleaner with potassium peroxymonosulphate	Environmental cultures once a week for patients who have undergone catheterization. Nurses working in wards were educated on infection control, contact precautions and hand hygiene	Reduction to one ESBL-producing strain 1 month later
Zhu et al. <sup>30</sup>	Thirty-one patients with <i>Klebsiella pneumoniae</i> carbapenemase-2 and/or New Delhi Metallo-β-Lactamase 1 <i>K. pneumoniae</i> strains in 10 different wards in a 2,076-bed tertiary care teaching hospital	Antimicrobial stewardship workers enhanced disinfection of the patient's room at least three times a day; after discharge, the surroundings were cleaned in depth using a 500-ppm chlorine solution and high-touch surfaces were emphasized by the hospital hygiene nurse manager for cleaning to reduce cross-transmission	Set up a working group on the antimicrobial stewardship. Strict infection control procedures were put in place, including rapid communication, suggestions on rational drug use in time and training courses, contact precautions of carriers, surveillance cultures of the hand and environment and screening for all contact patients	Return to the epidemiological stage with sporadic occurrence of carbapenem-resistant <i>K. pneumoniae</i> strains
Gavaldá et al. <sup>31</sup>	Increase in new cases of extensively drug-resistant (XDR) <i>Acinetobacter baumannii</i> after a period of high endemicity	To agree on the specific responsibilities and allocation of cleaning duties of the auxiliary nurses and housekeepers; deep environmental cleaning after a staggered schedule; increasing the cleaning frequency of high-touch surfaces; apply ready-to-use disinfectant wipes with cationic surfactant tensioactives, quaternary ammonium compounds and polymeric biguanide by auxiliary nurses and microfibre cleaning system containing 0.1% chlorine solution by housekeepers with the one-room, one-wipe approach	A multidisciplinary task force was established to determine the main control strategies; rectal swabs from patients admitted to the intensive care units	The implementation of interventions was associated with a significant change in the level of -8,360 XDR <i>A. baumannii</i> -infected patients per 1,000 patient days
Grabsch et al. <sup>32</sup>	Markedly increased colonized patients on active screening of 'high-risk' wards (intensive care unit and renal, liver transplant and haematology-oncology units) and VRE bacteraemia	Replacement of surface cleaning/disinfection compounds with sodium hypochlorite 1,000 ppm + detergent; employment of cleaning supervisors and establishment of a formal training programme for cleaning staff; implementation of a thrice-annual schedule detailed bleach-based 'super-clean-disinfection'	Adherence to the Centers for Disease Control and Prevention VRE guidelines for managing patients with colonization	Significant reductions in newly recognized colonized VRE (208/1,948 patients screened versus 324/4,035, a 24.8% reduction); the total burden of inpatients with VRE declined; median percentage of colonized inpatients per week: 19.4% versus 17.3%; hospital-wide VRE bacteraemia declined 83.1%
Salazar et al. <sup>33</sup>	Increase in CRE strains in a general hospital	Training to cleaning and disinfection staff twice a year; monthly supervisions of the cleaning and disinfection process with a checklist; reviewing for the cleanliness of surfaces with alcohol wipes, and analysed the processes	Full management support for all decisions related to infection control; blinded observation of hand hygiene with monthly evaluations, disseminating the data to sensitize health personnel; restrictive antimicrobial stewardship; active surveillance of healthcare infections and contact isolation of carries in a specific area	Decrease in the CRE rate: 0.59 versus 0.18 cases/1,000 patient days

CRE = carbapenem-resistant Enterobacteriaceae; ESBL = extended-spectrum beta-lactamase; VRE = vancomycin-resistant enterococci; XDR = extensively drug-resistant.

## Point 2: Assessing the hospital resistance profile

This involves identifying the most frequent MDR microorganisms and their distribution. In 2017, the World Health Organization published a list of critical priority pathogens. The most significant in hospitals include methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant enterococci, extended-spectrum beta-lactamase (ESBL) Enterobacteriaceae, Enterobacterales, *Clostridioides difficile* (CD), carbapenem-resistant *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and Enterobacteriaceae.<sup>2,12</sup> The distribution of MDR microorganisms varies in each hospital, region and continent.<sup>13,14</sup> Identifying the most common MDR microorganisms in a hospital, the areas where they arise and their behaviour over time helps focus efforts on areas needing surveillance and control.

## Point 3: Analysing antibiotic consumption

This includes evaluating antibiotic consumption with recommended measures.<sup>15,16</sup> These data and the epidemiological profile help determine if there are alarming antibiotic consumption patterns, such as frequent carbapenem use in an intensive care unit, with only 3% ESBL isolates. Antimicrobial stewardship includes electronic pre-authorization systems or personnel providing feedback after prescriptions; this can encompass all or some antimicrobials and all or certain hospital areas. Involving each department's staff in decision-making, protocol development and educational measures on prescribing is essential to achieving the main programme goal: changing antibiotic prescribing behaviour.<sup>15,17</sup>

Biochemist pharmacists should be included to report inappropriate antibiotic use and potentially lead the antimicrobial stewardship programme.<sup>18–20</sup> The microbiology laboratory should standardize culture reports to guide physicians towards appropriate treatment decisions.<sup>21–23</sup> Initially, programme goals should be defined, such as reducing CD rates or glycopeptide consumption, and compliance should be re-evaluated.

All available tools should be used to control antibiotics in the hospital through various interventions to improve antibiotic prescriptions.

## Point 4: Evaluating the cleaning and disinfection process

None of the previous points will be effective without environmental control. Microorganisms can live on surfaces for weeks or months, potentially causing infections in hospitalized patients through horizontal transmission.<sup>24,25</sup> Studies have shown that the risk of acquiring an MDR microorganism in a room previously occupied by a colonized patient is up to four times higher.<sup>26–28</sup> Different cleaning and disinfection interventions have controlled MDR microorganism outbreaks, highlighting the importance of this process (see *Table 1*).<sup>29–33</sup>

First, the process should be observed: how it is conducted and if basic principles and recommended frequencies are applied with correct materials and dilutions.<sup>34,35</sup> Disinfectants should meet international

standards, and selection should be based on local epidemiology.<sup>28,30</sup> Possible barriers, such as toxicity data, difficult dilutions, application methods and communication challenges that limit process compliance, should also be evaluated.<sup>36,37</sup> For well-resourced institutions, non-touch technologies are advantageous.<sup>38–40</sup> However, hospitals with fewer resources must consider human and material limitations. Frequent supervision of the process is also crucial: both observational and objective with methods, such as adenosine triphosphate bioluminescence, along with feedback and periodic update training.<sup>36,41,42</sup>

## Point 5: Promoting hand hygiene

The importance of adherence to hand hygiene in infection control is undeniable.<sup>43–45</sup> Every hospital should monitor hand hygiene adherence, disseminate these data and promote it. There is electronic technology that monitors product use and/or free electronic tools that consolidate hand hygiene observation data.<sup>46,47</sup>

## Point 6: Other measures

In certain contexts, the contact isolation of patients colonized or infected with MDR microorganisms is questionable, but it remains a recommended measure by the Centers for Disease Control and Prevention.<sup>48–50</sup> Contact isolation should be applied according to the epidemiological profile and hospital resources, educating staff on the use of personal protective equipment.

Other measures influencing MDR identification include protocols for detecting colonized patients, the use of skin-decolonizing agents and the emergence and study of outbreaks.<sup>51</sup> Equally important is having sufficient, permanent and trained staff for patient care, especially in critical areas, to prevent horizontal transmission, as well as infrastructure that allows individual patient rooms in some hospitals versus shared spaces where contamination is more likely.<sup>52–54</sup>

Although the importance of MDR control is globally recognized, not all countries have mandatory regulatory standards for hospital control. It is essential to know local regulations that support infection control interventions and request financial, human and technological support from executive directive management, considering cost–benefit and savings analyses.<sup>55–57</sup>

## Conclusion

MDR is a global threat, and its control is vital in hospitals. This control must be achieved through multiple measures (stewardship programme, improved cleaning and disinfection, adherence to hand hygiene, contact isolation and detection of colonized patients, among others) adopted after an epidemiological analysis that identifies frequent microorganisms and infections in hospital and through a working group that includes every actor from executive directive management to cleaning and disinfection staff, all fulfilling defined roles in evaluating, educating and continuously providing feedback on all MDR control processes in the hospital. □

1. Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *Lancet*. 2022;399:629–55. DOI: 10.1016/S0140-6736(21)02724-0.
2. Siegel J, Rhinehart E, Jackson M, et al. Management of multidrug-resistant organisms in healthcare settings, 2006. 2006. Available at: [www.cdc.gov/infection-control/hcp/mdro-management/index.html](http://www.cdc.gov/infection-control/hcp/mdro-management/index.html) (Date last accessed: 15 February 2017).
3. Cassini A, Högberg LD, Plachouras D, Burden of AMR Collaborative Group. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015:

A population-level modelling analysis. *Lancet Infect Dis*. 2019;19:56–66. DOI: 10.1016/S1473-3099(18)30605-4.

4. Zimmerman FS, Assous MV, Bdolah-Abram T, et al. Duration of carriage of carbapenem-resistant Enterobacteriaceae following hospital discharge. *Am J Infect Control*. 2013;41:190–4. DOI: 10.1016/j.ajic.2012.09.020.
5. Riccio ME, Verschuuren T, Conzelmann N, et al. Household acquisition and transmission of extended-spectrum B-lactamase (ESBL)-producing enterobacteriaceae after hospital discharge of ESBL-positive index patients. *Clin Microbiol Infect*. 2021;27:1322–9. DOI: 10.1016/j.cmi.2020.12.024.

6. Gidey K, Gidey MT, Hailu BY, et al. Clinical and economic burden of healthcare-associated infections: A prospective cohort study. *PLoS One*. 2023;18:e0282141. DOI: 10.1371/journal.pone.0282141.
7. National Healthcare Safety Network. Bloodstream infection event (central line-associated bloodstream infection and non-central line associated bloodstream infection) 2024. Available at: [www.cdc.gov/nhsn/pdfs/pscmanual/4psc\\_clabscurrent.pdf](http://www.cdc.gov/nhsn/pdfs/pscmanual/4psc_clabscurrent.pdf) (Date last accessed: 8 August 2024).
8. National Healthcare Safety Network. Pneumonia (ventilator-associated [VAP] and non-ventilator-associated pneumonia [PNEU] event) 2024. Available at: [www.cdc.gov/nhsn/pdfs/](http://www.cdc.gov/nhsn/pdfs/)

- psmanual/6psvcapcurrent.pdf (Date last accessed: 8 August 2024).
- Wang M, Wei H, Zhao Y, et al. Analysis of multidrug-resistant bacteria in 3223 patients with hospital-acquired infections (HAI) from a tertiary general hospital in China. *Bosn J Basic Med Sci.* 2019;19:86–93. DOI: 10.17305/bjbm.2018.3826.
  - Cornejo-Juárez P, Vilar-Compte D, Pérez-Jiménez C, et al. The impact of hospital-acquired infections with multidrug-resistant bacteria in an oncology intensive care unit. *Int J Infect Dis.* 2015;31:31–4. DOI: 10.1016/j.ijid.2014.12.022.
  - Jiang A-M, Shi X, Liu N, et al. Nosocomial infections due to multidrug-resistant bacteria in cancer patients: A six-year retrospective study of an oncology center in Western China. *BMC Infect Dis.* 2020;20:452. DOI: 10.1186/s12879-020-05181-6.
  - World Health Organization (WHO). Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug-resistant bacterial infections, including tuberculosis 2017. Available at: [www.who.int/publications/i/item/WHO-EMP-IAU-2017.12](http://www.who.int/publications/i/item/WHO-EMP-IAU-2017.12) (Date last accessed: 8 August 2024).
  - European Centre for Disease Prevention and Control and World Health Organization European Region. Antimicrobial resistance surveillance in Europe 2023 – 2021 data. 2023. Available at: [www.ecdc.europa.eu/sites/default/files/documents/Antimicrobial%20resistance%20surveillance%20in%20Europe%202023%20-%202021%20data.pdf](http://www.ecdc.europa.eu/sites/default/files/documents/Antimicrobial%20resistance%20surveillance%20in%20Europe%202023%20-%202021%20data.pdf) (Date last accessed: 8 August 2024).
  - Seas C, Garcia C, Salles MJ, et al. Latin America Working Group on Bacterial Resistance. *Staphylococcus aureus* bloodstream infections in Latin America: Results of a multinational prospective cohort study. *J Antimicrob Chemother.* 2018;73:212–22. DOI: 10.1093/ajic/dkx350.
  - World Health Organization. Antimicrobial stewardship programmes in health-care facilities in low- and middle-income countries. A WHO Practical Toolkit. 2019. Available at: <https://iris.who.int/bitstream/handle/10665/329404/9789241515481-eng.pdf?sequence=1> (Date last accessed: 8 August 2024).
  - Barlam TF, Cosgrove SE, Abbo LM, et al. Implementing an antibiotic stewardship program: Guidelines by the Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America. *Clin Infect Dis.* 2016;62:e51–77. DOI: 10.1093/cid/ciw118.
  - WHO Regional Office for Europe. Antimicrobial stewardship interventions: A practical guide. 2021. Available at: <https://iris.who.int/bitstream/handle/10665/340709/9789289054980-eng.pdf> (Date last accessed: 8 August 2024).
  - Cantudo-Cuenca MR, Jiménez-Morales A, Martínez-de la Plata JE. Pharmacist-led antimicrobial stewardship programme in a small hospital without infectious diseases physicians. *Sci Rep.* 2022;12:9501. DOI: 10.1038/s41598-022-13246-6.
  - Abdelrahman DH, AbuSara AK, Khabour DS. The impact of pharmacist-led antimicrobial stewardship review of cultures in the ambulatory setting at a comprehensive cancer center. *Hosp Pharm.* 2023;58:392–5. DOI: 10.1177/00185787221150920.
  - Jantarathaneewat K, Camins B, Apisarnthanarak A. The role of the clinical pharmacist in antimicrobial stewardship in Asia: A review. *Antimicrob Steward Healthc Epidemiol.* 2022;2:e176. DOI: 10.1017/ash.2022.310.
  - Coupat C, Pradier C, Degand N, et al. Selective reporting of antibiotic susceptibility data improves the appropriateness of intended antibiotic prescriptions in urinary tract infections: A case-vignette randomised study. *Eur J Clin Microbiol Infect Dis.* 2013;32:627–36. DOI: 10.1007/s10096-012-1786-4.
  - Wang Y, Zhang X, Zhou Q, Xu X. Impact of selective reporting of antimicrobial susceptibility testing report on clinicians' prescribing behavior of antibiotics. *Front Pharmacol.* 2023;14:1225531. DOI: 10.3389/fphar.2023.1225531.
  - Lestin-Bernstein F, Harberg R, Schumacher I, et al. *Staphylococcus aureus* – Selective reporting of antibiogram results and its impact on antibiotic use: Interventional study with a reference group on the effect of switching from non-selective to selective antibiotic reporting. *Antimicrob Resist Infect Control.* 2021;10:157. DOI: 10.1186/s13756-021-01021-7.
  - Porter L, Sultan O, Mitchell BG, et al. How long do nosocomial pathogens persist on inanimate surfaces? A scoping review. *J Hosp Infect.* 2024;147:25–31. DOI: 10.1016/j.jhin.2024.01.023.
  - Sood G, Perl TM. Outbreaks in health care settings. *Infect Dis Clin North Am.* 2016;30:661–87. DOI: 10.1016/j.idc.2016.04.003.
  - Drees M, Snyderman DR, Schmid CH, et al. Prior environmental contamination increases the risk of acquisition of vancomycin-resistant *Enterococci*. *Clin Infect Dis.* 2008;46:678–85. DOI: 10.1086/527394.
  - Nseir S, Blazejewski C, Lubret R, et al. Risk of acquiring multidrug-resistant gram-negative Bacilli from prior room occupants in the intensive care unit. *Clin Microbiol Infect.* 2011;17:1201–8. DOI: 10.1111/j.1469-0691.2010.03420.x.
  - Shaughnessy MK, Miceli RL, DePestel DD, et al. Evaluation of hospital room assignment and acquisition of *Clostridium difficile* infection. *Infect Control Hosp Epidemiol.* 2011;32:201–6. DOI: 10.1086/658669.
  - Ikeda Y, Shigemura K, Nomi M, et al. Infection control following an outbreak of extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae* isolated from catheter-associated urinary tract infection. *Jpn J Infect Dis.* 2018;71:158–61. DOI: 10.7883/yoken.JJID.2017.330.
  - Zhu J, Li Q, Li X, et al. Successful control of the first carbapenem-resistant *Klebsiella pneumoniae* outbreak in a Chinese hospital 2017–2019. *Antimicrob Resist Infect Control.* 2020;9:91. DOI: 10.1186/s13756-020-00757-y.
  - Gavaldà L, Soriano AM, Cámara J, et al. Control of endemic extensively drug-resistant *Acinetobacter baumannii* with a cohorting policy and cleaning procedures based on the 1 room, 1 wipe approach. *Am J Infect Control.* 2016;44:520–4. DOI: 10.1016/j.ajic.2015.11.036.
  - Grabsch EA, Mahony AA, Cameron DRM, et al. Significant reduction in vancomycin-resistant *Enterococcus* colonization and *Bacteraemia* after introduction of a bleach-based cleaning-disinfection programme. *J Hosp Infect.* 2012;82:234–42. DOI: 10.1016/j.jhin.2012.08.010.
  - Salazar Tamayo GJ, Castelo Vigme JP, Peralta Prado AB, et al. How to avoid multidrug resistance (MDR)? Successful experience with the application of a hospital bundle (HB). Presented at: ECCMID 2023, Copenhagen, Denmark Available at: <https://elibrary.escmid.org/?search%5Bfrom%5D=0&search%5Bquery%5D=%22Grace%22&search%5Bcategory%5D%5B0%5D=8&search%5Bdate%5D=&tab=docs#results>.
  - Rutala WA, Weber DJ. Healthcare Infection Control Practices Advisory Committee (HICPAC). Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008. Available at: [www.cdc.gov/infection-control/media/pdfs/Guideline-Disinfection-H.pdf](http://www.cdc.gov/infection-control/media/pdfs/Guideline-Disinfection-H.pdf) (Date last accessed: 8 August 2024).
  - Rutala WA, Weber DJ. Disinfection and sterilization in health care facilities: What clinicians need to know. *Clin Infect Dis.* 2004;39:702–9. DOI: 10.1086/423182.
  - Assadian O, Harbarth S, Vos M, et al. Practical recommendations for routine cleaning and disinfection procedures in healthcare institutions: A narrative review. *J Hosp Infect.* 2021;113:104–14. DOI: 10.1016/j.jhin.2021.03.010.
  - Rutala WA, Weber DJ. Sterilization, high-level disinfection, and environmental cleaning. *Infect Dis Clin North Am.* 2011;25:45–76. DOI: 10.1016/j.idc.2010.11.009.
  - Dancer SI. Controlling hospital-acquired infection: Focus on the role of the environment and new technologies for decontamination. *Clin Microbiol Rev.* 2014;27:665–90. DOI: 10.1128/CMR.00020-14.
  - Scott R, Joshi LJ, McGinn C. Hospital surface disinfection using ultraviolet germicidal irradiation technology: A review. *Health Technol Lett.* 2022;9:25–33. DOI: 10.1049/htl2.12032.
  - Boyce JM. Modern technologies for improving cleaning and disinfection of environmental surfaces in hospitals. *Antimicrob Resist Infect Control.* 2016;5:10. DOI: 10.1186/s13756-016-0111-x.
  - Public Health Ontario. Best practices for environmental cleaning for prevention and control of infections in all health care settings, 3rd edition. 2018. Available at: [www.publichealthontario.ca/-/media/documents/B/2018/bp-environmental-cleaning.pdf](http://www.publichealthontario.ca/-/media/documents/B/2018/bp-environmental-cleaning.pdf) (Date last accessed: 8 August 2024).
  - Smith PW, Beam E, Sayles H, et al. Impact of adenosine triphosphate detection and feedback on hospital room cleaning. *Infect Control Hosp Epidemiol.* 2014;35:564–9. DOI: 10.1086/675839.
  - Pittet D, Hugonnet S, Harbarth S, et al. Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. Infection control programme. *Lancet.* 2000;356:1307–12. DOI: 10.1016/S0140-6736(00)02814-2.
  - Kingston L, O'Connell NH, Dunne CP. Hand hygiene-related clinical trials reported since 2010: A systematic review. *J Hosp Infect.* 2016;92:309–20. DOI: 10.1016/j.jhin.2015.11.012.
  - Marimuthu K, Pittet D, Harbarth S. The effect of improved hand hygiene on nosocomial MRSA control. *Antimicrob Resist Infect Control.* 2014;3:34. DOI: 10.1186/2047-2994-3-34.
  - Allegranzi B, Gayet-Ageron A, Damani N, et al. Global implementation of WHO's multimodal strategy for improvement of hand hygiene: A quasi-experimental study. *Lancet Infect Dis.* 2013;13:843–51. DOI: 10.1016/S1473-3099(13)70163-4.
  - Wang C, Jiang W, Yang K, et al. Electronic monitoring systems for hand hygiene: Systematic review of technology. *J Med Internet Res.* 2021;23:e27880. DOI: 10.2196/27880.
  - Kelly G, Hudson M, Apple B, et al. Discontinuation of contact precautions in patients with hospital-acquired MRSA and VRE infections during the COVID-19 pandemic: A multi-center experience. *J Infect Prev.* 2024;25:33–7. DOI: 10.1177/17571774231208312.
  - Gottlieb LB, Walits E, Patel G, Schaefer S. Taking off the gown: Impact of discontinuing contact precautions for extended-spectrum  $\beta$ -lactamase (ESBL)-producing organisms. *Antimicrob Steward Healthc Epidemiol.* 2021;1:e31. DOI: 10.1017/ash.2021.189.
  - Siegel JD, Rhinehart E, Jackson M, et al. 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings Available at: [www.cdc.gov/infection-control/media/pdfs/Guideline-Isolation-H.pdf](http://www.cdc.gov/infection-control/media/pdfs/Guideline-Isolation-H.pdf) (Date last accessed: 8 August 2024).
  - Tien KL, Wang JT, Sheng WH, et al. Chlorhexidine bathing to prevent healthcare-associated vancomycin-resistant *Enterococcus* infections: A cluster quasi-experimental controlled study at intensive care units. *J Formos Med Assoc.* 2021;120:1014–21. DOI: 10.1016/j.jfma.2020.08.048.
  - Stone PW, Pogorzelska M, Kunches L, et al. Hospital staffing and health care-associated infections: A systematic review of the literature. *Clin Infect Dis.* 2008;47:937–44. DOI: 10.1086/591696.
  - Wang R, Xia J, Zhao S, et al. The advantages of single-room management in an ICU and the changing trend of drug-resistant bacteria over the last 5 years. *Intensive Care Res.* 2023;3:18–29. DOI: 10.1007/s44231-023-00028-5.
  - Halaby T, Al Naiemi N, Beishuizen B, et al. Impact of single room design on the spread of multi-drug resistant bacteria in an intensive care unit. *Antimicrob Resist Infect Control.* 2017;6:117. DOI: 10.1186/s13756-017-0275-z.
  - Bryant KA, Harris AD, Gould CV, et al. Necessary infrastructure of infection prevention and healthcare epidemiology programs: A review. *Infect Control Hosp Epidemiol.* 2016;37:371–80. DOI: 10.1017/ice.2015.333.
  - Benenson S, Cohen MJ, Schwartz C, et al. Is it financially beneficial for hospitals to prevent nosocomial infections? *BMC Health Serv Res.* 2020;20:653. DOI: 10.1186/s12913-020-05428-7.
  - Xiong CL, Wang GG, Hanafi WU. Financial impact of nosocomial infections on surgical patients in an eastern Chinese hospital: A propensity score matching study. *J Hosp Infect.* 2023;139:67–73. DOI: 10.1016/j.jhin.2023.05.017.